



Postoperative Instructions: Quad/Patellar Tendon Repair

Please read these instructions thoroughly. Most of your questions will be addressed here. The first page is a summary. You can find additional details on the following pages. Following these precautions will help minimize complications and address concerns about your surgery. Additional information can be found at www.bradleysmithmd.com.

SUMMARY PAGE

Activity: You may be weight-bearing as tolerated with the knee brace locked straight.

Brace: Wear the brace at all times except to remove for hygiene purposes or if you are doing exercises with physical therapy. It should be locked straight at all times until directed otherwise.

Medications:

Oxycodone 5 mg: 1-2 tabs by mouth every 4-6 hours as needed for severe pain.

Methocarbamol 750 mg (optional): 1 tab by mouth every 8-12 hours as needed for muscle spasms.

Tylenol 325 mg: we recommend 2 tabs (650 mg) every 6 hours until your pain is low enough for you to switch to taking this as needed.

Aleve 220 mg (Naproxen): we recommend 2 tabs every 12 hours for the first 5 days after surgery.


Aspirin 81 mg: 1 tab two times per day to reduce the risk of blood clots.

Pain control: You likely received a nerve block for postoperative pain control. We recommend that you start your pain medication when you get home from the hospital. Continue the medication for the first 1-2 days until the block wears off and you have a better idea of how you will handle the pain. Keep your leg **elevated** and use **ice** as much as possible to help with pain and swelling.

Dressing: You may remove the outer ACE wrap and bandage 1-2 days after surgery. There is a waterproof dressing underneath that you should leave until your followup appointment.

myHealthTrack: Quad/Patellar Tendon Repair

Follow up: 10-14 days



Office: (801) 747-1020

Fax: (801) 747-1023

www.bradleysmithmd.com



Pain Control and Medications

You may have received a nerve block or a local anesthetic injection to help with pain control. This will likely wear off 8-12 hours after surgery. You should start taking pain medication when you get home and before you go to sleep so that you have something working before the block wears off. Otherwise the pain can be difficult to control. Typically, pain is the worst during the first 2-3 days after surgery before starting to subside.

Medications:

Prescription:

- Oxycodone 5 mg: this is a strong narcotic medication. You may take up to 1-2 tablets every 4-6 hours AS NEEDED for the first few days after surgery. After 2-4 days, you should be able to space out or discontinue this medication. Do not drive or drink alcohol while taking this medication.
- Methocarbamol 750 mg: this is a muscle-relaxer that can help with muscle spasms and pain. Be careful taking this medication with narcotic pain medications, as they both can cause drowsiness. Do not drive or operate heavy machinery while taking.

Over-the-counter:

- Tylenol (Acetaminophen) 325 mg: this is a strong non-narcotic medication for pain. You should take 2 tablets (650 mg) every 6 hours. This can be taken with the oxycodone, and can reduce the amount of oxycodone that you need to reduce your pain.
- Aleve (Naproxen) 220 mg: this is a strong anti-inflammatory medication. We recommend that you take 2 tablets (440 mg) every 12 hours with meals for the first 5 days after surgery. Do **NOT** take this medication if you have a history of gastric ulcers or sensitivity to other anti-inflammatory medications.
- Aspirin 81 mg: this medication is to help prevent blood clots after surgery. Please take 81 mg two times daily for a total of 2 weeks after surgery.

If you have any side effects (i.e. nausea, rash, trouble breathing) from a medication, discontinue its use and call our office. Constipation is common after surgery and while you are taking narcotics for pain control. Drink plenty of clear liquids, and consider using an over-the-counter stool softener such as Docusate or Miralax.

Wound Care and Showering

- It is normal for there to be bleeding and swelling following surgery – if blood saturates the dressing, simply reinforce with additional gauze dressing for the remainder of day and check again.



Office: (801) 747-1020

Fax: (801) 747-1023


www.bradleysmithmd.com



- You may remove the outer ACE wrap and bandage 1-2 days after surgery. Underneath you will find a mesh tape with dermabond glue. Leave this in place until your follow up appointment.
- You may shower with the mesh tape/glue dressing.
- Do not soak the incision in water. Do not go swimming in the pool or lake/ocean until Dr. Smith tells you that you can do so.
- Please do not use Bacitracin or other ointments under the bandage.

Physical Therapy

- Physical therapy is necessary after your surgery. You should plan to start physical therapy within 1-2 weeks of your surgery, and you may begin within a few days if you would like.
- An optional but beneficial supplement to physical therapy is the myHealthTrack app. This app can be downloaded using the QR code or by going to mht.link/utahortho. Then choose **Dr. Smith → I'm Having / I've Had Surgery → Knee → Quad/Patellar Tendon Repair**. Make sure to clarify which track you should follow before starting.



Activity

- You may weight-bear as tolerated on the operative leg with the knee brace locked straight (extension).
- **Crutches** are recommended for the first 5-7 days to help provide support and aid in ambulation, and then as needed after that.
- The knee brace needs to be locked straight at all times. The brace also must be worn while sleeping at night. You may remove the brace for hygiene purposes, and when doing exercises with physical therapy.

Ice and Elevation

- When sitting or lying down be sure to keep your leg elevated up above the level of your heart to help decrease swelling.
- Ice should be used as much as possible for the first 3-5 days to help decrease swelling.
- After the first 3-5 days, ice your knee 4-5 times per day as needed.
- Never place ice directly on skin. Use a thin towel or clothing between the ice and your skin.

Follow-Up

- Please contact the office to schedule an appointment 10-14 days after your surgery.



Office: (801) 747-1020

Fax: (801) 747-1023

www.bradleysmithmd.com



- You will be seen by Dr. Smith for your first postoperative visit to review your intraoperative findings and to go over any questions you may have.

Driving

- Do not drive a car or operate machinery while taking narcotic pain medication.
- Driving is not recommended for 6 weeks following surgery. After that point, you can return to driving when you are able to walk comfortably without assistive devices. If surgery was on your left leg and you

When to call for help

Call 911 anytime you think you may need emergency care. For example, call if:

- You pass out (lose consciousness).
- You have severe trouble breathing.
- You have sudden chest pain and shortness of breath, or you cough up blood.
- You have severe nausea or vomiting.

Call the office if:

- You have pain that does not go away after you take pain pills.
- You have a fever over 101.3°F.
- You have loose stitches, or your incision comes open.
- Your incision keeps bleeding 3 days after your surgery.
- You have signs of infection, such as redness around the incision or pus draining from your incision.



Office: (801) 747-1020

Fax: (801) 747-1023

www.bradleysmithmd.com