

## Postoperative Instructions: Basic Knee Arthroscopy

This instruction packet is intended for patients undergoing simple knee arthroscopy surgery. If you had a ligament, tendon, cartilage, or meniscus repair/reconstruction, you may need the Knee Surgery Instructions instead. Please read these instructions thoroughly. Most of your questions will be addressed here. The first page is a summary. You can find additional details on the following pages. Following these precautions will help minimize complications and address concerns about your surgery. Additional information can be found at [bradleysmithmd.com](http://bradleysmithmd.com).

### SUMMARY PAGE

**Weight-bearing status:** Weight-bearing as tolerated

**Range of Motion:** Progress as tolerated

#### **Medications:**

**Oxycodone 5 mg:** 1 tab by mouth every 4-6 hours as needed for severe pain.

**Tylenol 325 mg:** we recommend 2 tabs (650 mg) every 6 hours until your pain is low enough for you to switch to taking this as needed.

**Aleve 220 mg (Naproxen):** we recommend 2 tabs every 12 hours for the first 5 days after surgery.


**Aspirin 81 mg:** 1 tab two times per day for 14 days to reduce the risk of blood clots.

#### **Pain control:**

We recommend that you start your pain medication when you get home from the hospital. You should take Tylenol and/or Aleve first, and only use the oxycodone when your pain is not completely controlled by the other medications. Keep your leg **elevated** and use **ice or a cryo-cuff** as much as possible to help with pain and swelling.

**myHealthTrack:** Knee Arthroscopy

**Follow up:** 10-14 days





## Pain Control and Medications

You may have received a nerve block or a local anesthetic injection to help with pain control. This will likely wear off 8-12 hours after surgery. You should start taking pain medication when you get home and before you go to sleep so that you have something working before the block wears off. Otherwise the pain can be difficult to control. Typically, pain is the worst during the first 2-3 days after surgery before starting to subside.

### **Medications:**

#### Prescription:

- Oxycodone 5 mg: this is a strong narcotic medication. You may take up to 1-2 tablets every 4-6 hours AS NEEDED for the first few days after surgery. After 2-4 days, you should be able to space out or discontinue this medication. Do not drive or drink alcohol while taking this medication.

#### Over-the-counter:

- Tylenol (Acetaminophen) 325 mg: this is a strong non-narcotic medication for pain. You should take 2 tablets (650 mg) every 6 hours. This can be taken with the oxycodone, and can reduce the amount of oxycodone that you need to reduce your pain.
- Aleve (Naproxen) 220 mg: this is a strong anti-inflammatory medication. We recommend that you take 2 tablets (440 mg) every 12 hours with meals for the first 5 days after surgery. You may stop taking this medication sooner if your pain is controlled. Do **NOT** take this medication if you have a history of gastric ulcers or sensitivity to other anti-inflammatory medications.
- Aspirin 81 mg: this medication is to help prevent blood clots after surgery. Please take 81 mg two times per day for a total of 3 weeks after surgery.

If you have any side effects (i.e. nausea, rash, trouble breathing) from a medication, discontinue its use and call our office. Constipation is common after surgery and while you are taking narcotics for pain control. Drink plenty of clear liquids, and consider using an over-the-counter stool softener such as Docusate or Miralax.



Office: (801) 747-1020

Fax: (801) 747-1023


[www.bradleysmithmd.com](http://www.bradleysmithmd.com)

## Wound Care and Showering

- Leave the surgical bandages on and do not shower for 48 HOURS.
- It is normal for the joint to bleed and swell following surgery – if blood soaks onto the ACE bandage, simply reinforce with additional gauze dressing for the remainder of day and check again.
- The ACE bandage can be laundered in normal fashion and reused.
- You may loosen the bandage if swelling or progressive numbness occurs in the extremity.
- After 48 hours, remove bandages and gauze. It is normal to see bloody fluid on the gauze sponges.
- If you have steri-strips, leave them in place over the incisions until your first appointment.
- After removing the dressing, you may shower. You can get the incisions wet in the shower and let water run over the incisions. Gently pat the incisions dry when done, and cover the incisions with band-aids.
- Do NOT use Bacitracin, lotions, or other ointments to the incisions.
- Do NOT soak incisions in any pool/bathwater until 3 weeks postoperatively or until your incisions are completely closed.
- Do not allow pets to sit on your lap or sleep in your bed for at least 4 weeks following surgery. Pets may harbor flies, mites, or other organisms that may cause an infection.

## Physical Therapy

- Physical therapy is not always necessary after arthroscopic knee surgery. The need for physical therapy will be discussed at your first postoperative appointment.
- You may progress range of motion as tolerate
- There are home exercises on the last page that you can begin as soon as you feel comfortable after surgery.
- An optional but beneficial supplement to physical therapy is the myHealthTrack app. This app can be downloaded using the QR code or by going to [mht.link/utahortho](http://mht.link/utahortho). Then choose **Dr. Smith** → **Operative** → **Knee** → **Knee Arthroscopy**.



## Activity

- Weight-bearing after surgery depends on the type of procedure you have performed. After most routine knee arthroscopies, you can weight-bear as tolerated. Make sure to clarify your weight-bearing status with your nurse or Dr. Smith prior to discharge from the surgery center.
- **Crutches** are recommended for the first 3-5 days to help provide support and aid in ambulation.



## Ice and Elevation

- Try and rest the first few days following surgery.
- When sitting or lying down be sure to keep your leg elevated up above the level of your heart to help decrease swelling.
- Ice should be used as much as possible for the first 3-5 days to help decrease swelling.
- After the first 3-5 days, ice your knee 4-5 times per day as needed.
- Never place ice directly on skin. Use a thin towel or clothing between the ice and your skin.

## Follow-Up

- You have likely already been scheduled for your first postoperative visit.
- If you have not received an appointment please contact the office to schedule an appointment 10-14 days after your surgery.
- You will be seen by Dr. Smith for your first postoperative visit to review your intraoperative findings and to go over any questions you may have. Sutures will also be removed at that time.

## Driving

- Do not drive a car or operate machinery while taking narcotic pain medication.
- In general, you can return to driving when you are able to walk comfortably without assistive devices.

## When to call for help

**Call 911 anytime you think you may need emergency care. For example, call if:**

- You pass out (lose consciousness).
- You have severe trouble breathing.
- You have sudden chest pain and shortness of breath, or you cough up blood.
- You have severe nausea or vomiting.

**Call the office if:**

- You have pain that does not go away after you take pain pills.
- You have a fever over 101.3 °F.
- You have loose stitches, or your incision comes open.
- Your incision keeps bleeding 3 days after your surgery.
- You have signs of infection, such as redness around the incision or pus draining from your incision.



Office: (801) 747-1020

Fax: (801) 747-1023

[www.bradleysmithmd.com](http://www.bradleysmithmd.com)




**BRADLEY W SMITH MD**

Sports Medicine Orthopedic Surgeon

**YOU DESERVE A COMEBACK STORY**

## Home Knee Exercises

### QUAD SETS




Tighten muscles on top of thigh by pushing knee down to floor or table.

Hold 1-2 seconds. Repeat 10-15 times

Do 3 sessions per day.

### STRAIGHT LEG RAISE




Tighten muscle on front of thigh then lift leg 8-10 inches from floor keeping knee locked.

Hold 1-2 seconds. Repeat 10-15 times

Do 3 sessions per day.

### CALF PUMPS



Relax leg. Gently bend and straighten ankle. Move through full range of motion. Avoid pain.

Hold 1-2 seconds. Repeat 10-15 times

Do 3 sessions per day

### LEG HANGS WITH GRAVITY




Allow leg to hang with gravity and bend as tolerated for 60-90 seconds.

DO NOT ACTIVELY STRAIGHTEN LEG FROM THE BENT POSITION

Do 3 times per day

### Heel Slides



Lie on your back with your legs straight and your feet slightly apart. Slide your affected leg as close to your buttocks as you can. Hold this position for 5 seconds. Slide your heel back to the starting position.

Do 1-3 sets of 10 repetitions, with 1 minute of rest between.

Try to do this 2-3 times per day